THE LANCET Global Health

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: Kiarie H, Temmerman M, Nyamai M, et al. The COVID-19 pandemic and disruptions to essential health services in Kenya: a retrospective time-series analysis. *Lancet Glob Health* 2022; **10:** e1257–67.

Appendix Table 1: Indicators of utilisation and access to essential health services

Category	Indicator	Target group
Access/Utilization of primary health care facilities	Outpatient visits	Adults, <5 years
Reproductive, maternal, newborn, child, and adolescent health (RMNCAH) Indicators	Skilled deliveries	15-49 years
	Antenatal care (ANC)	<5 years
	Children presenting with pneumonia	<5 years
	Vitamin A supplements	15-59 months
	DPT3 doses administered	<5 years
	Children attending Child Welfare Clinic (CWC) who are underweight	<5 years
	Rate of children treated for diarrhoea	<5 years
Sexual Violence	Sexual violence cases	All
Communicable Diseases	HIV tests conducted	All
	Persons with HIV infection newly started on ART	All
	TB number of notified cases	All
	Patients tested for malaria	All
	Confirmed cases of malaria	All
Non-Communicable Diseases	Cervical cancer screening	Women 25-49 years
	Hypertension cases	All
	Diabetes cases	All



